German Research Products - GRP GmbH
In der Stockwiese 26
D-85410 Haag/Amper, Germany
Email: info@grp-ak.de
Phone: +49 (0)8167 6335

Product Datasheet

Goat anti-Mouse IgG (H&L), F(ab)'2 fragment, TRITC conjugated, min. cross-reactivity to bovine, goat, human, rabbit, or rat IgG

GRP13095

Species/Host	Goat
Reactivity	Mouse
Tested Applications	FC, IL
Immunogen	Purified Mouse IgG, whole molecule
Form/Appearance	Lyophilized
Storage	Store at 2-8 °C.
Note	For research use only.
Clonality	Polyclonal
Purity	Affinity purified
Dilution Range	1 : 20-1 : 2000 (Flow cyt), (IL)
Application Notes	Additional Information: Antibody purity is > 90% based on SDS-PAGE Small amounts of intact IgG may be present.Affinity purified using solid phase Mouse IgG .Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free.Based on IEP, this antibody reacts with: • heavy (γ) chains on mouse IgG • light chains on all mouse immunoglobulins .Based on IEP, no reactivity is observed to: • non-immunoglobulin human serum immunoglobulins • bovine, goat, human, rabbit or rat IgG . Background: Goat anti-Mouse IgG (H&L), F(ab)'2 fragment, TRITC conjugated, min. cross-reactivity to bovine, goat, human, rabbit, or rat IgG is a secondary antibody conjugated to TRITC, which binds to mouse IgG (H&L), F(ab)'2 Fragment in immunological assays. Reconstitution: For reconstitution add 0.55 ml of sterile water.Let it stand 30 minutes at room temperature to dissolve. Prepare fresh working dilutions daily.