

## Product Datasheet

### Goat anti-Mouse IgM (H&L), TRITC conjugated GRP13121

|                     |                                    |
|---------------------|------------------------------------|
| Species/Host        | Goat                               |
| Reactivity          | Mouse                              |
| Tested Applications | FC, IL                             |
| Immunogen           | Purified Mouse IgM, whole molecule |
| Form/Appearance     | Lyophilized                        |
| Storage             | Store at 2-8 °C.                   |
| Note                | For research use only.             |
| Clonality           | Polyclonal                         |
| Purity              | Affinity purified                  |
| Dilution Range      | 1 : 20-1 : 2000 (Flow cyt), (IL)   |

**Application Notes** Additional Information: Antibody purity is > 95% based on SDS-PAGE. Affinity purified using solid phase Mouse IgM. Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/IgG free. 0.05% (w/v) Sodium Azide is added as a preservative. Based on IEP, this antibody reacts with: • heavy (γ) chains on mouse IgM • light chains on all mouse immunoglobulins. Based on IEP, no reactivity is observed to: • non-immunoglobulin mouse serum proteins. Background: Goat anti-Mouse IgM (H&L), TRITC conjugated is a secondary antibody conjugated to TRITC, which binds to mouse IgM in immunological assays. Reconstitution: For reconstitution add 1.1 ml of sterile water. Let it stand 30 minutes at room temperature to dissolve. Prepare fresh working dilutions daily.