

## Product Datasheet

### Goat anti-Rabbit IgG (H&L), min. cross-reactivity to bovine, goat, human, mouse or rat IgG or serum GRP13128

Species/Host	Goat
Reactivity	Rabbit
Immunogen	Purified Rabbit IgG, whole molecule
Form/Appearance	Liquid
Storage	Store at 2-8 °C.
Note	For research use only.
Clonality	Polyclonal
Purity	Affinity purified goat IgG, using solid phase rabbit IgG
Dilution Range	The optimal working dilution should be determined by the investigator.
Application Notes	<p>Additional Information: Antibody purity is &gt; 95% based on SDS-PAGE. Affinity purified using solid phase Rabbit IgG. Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2. 0.05% (w/v) Sodium Azide is added as a preservative. Based on IEP, this antibody reacts with:</p> <ul style="list-style-type: none"><li>• heavy (γ) chains on rabbit IgG</li><li>• light chains on all rabbit immunoglobulins</li></ul> <p>Based on IEP, no reactivity is observed to:</p> <ul style="list-style-type: none"><li>• non-immunoglobulin rabbit serum proteins</li><li>• serum proteins from bovine, goat, human, mouse, or rat</li><li>• IgG from bovine, goat, human, mouse or rat</li></ul> <p>Background: Goat anti-Rabbit IgG (H&amp;L), min. cross-reactivity to bovine, goat, human, mouse or rat IgG or serum is a secondary antibody which binds to rabbit IgG in immunological assays.</p>