

Product Datasheet

Donkey anti-Rabbit IgG (H&L), F(ab)'2 Fragment, min. cross-reactivity to Bovine, Chicken, Goat, Guinea Pig, Hamster, Horse, Human, Mouse, Rat or Sheep GRP13130

| | |
|-------------------|---|
| Species/Host | Donkey |
| Reactivity | Rabbit |
| Immunogen | Purified Rabbit IgG, whole molecule |
| Form/Appearance | Liquid |
| Storage | Store at 2-8 °C. |
| Note | For research use only. |
| Clonality | Polyclonal |
| Purity | Affinity purified |
| Dilution Range | The optimal working dilution should be determined by the investigator. |
| Application Notes | <p>Additional Information: Antibody purity is > 90% based on SDS-PAGE. Small amounts of intact IgG may be present. Affinity purified using solid phase Rabbit IgG. Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2. 0.05% (w/v) Sodium Azide is added as a preservative. Based on IEP, this antibody reacts with:</p> <ul style="list-style-type: none">• heavy (γ) chains on Rabbit IgG• light chains on all Rabbit immunoglobulins <p>Based on IEP, no reactivity is observed to:</p> <ul style="list-style-type: none">• non-immunoglobulin Rabbit serum immunoglobulins• IgG from Bovine, Chicken, Goat, Guinea Pig, Hamster, Horse, Human, Mouse, Rat or Sheep. <p>Background: Donkey anti-Rabbit IgG (H&L), F(ab)'2 Fragment, min. cross-reactivity to Bovine, Chicken, Goat, Guinea Pig, Hamster, Horse, Human, Mouse, Rat or Sheep is a secondary antibody which binds to rabbit IgG (H&L), F(ab)'2 Fragment in immunological assays.</p> |