

## Product Datasheet

### Goat anti-Rabbit IgG Fc, ALP conjugated, min. cross-reactivity to bovine horse or human serum proteins GRP13135

Species/Host	Goat
Reactivity	Rabbit
Tested Applications	FC, IL
Immunogen	Purified Rabbit IgG Fc
Form/Appearance	Liquid
Storage	Store liquid at Store at 2-8°C. For storage at -20 °C, dilute with an equal volume of glycerol to prevent loss of enzymatic activity. Prepare working dilutions prior to use and then discard.
Note	For research use only.
Clonality	Polyclonal
Purity	Affinity purified
Dilution Range	1 : 20-1 : 2000 (Flow cyt), (IL)
Application Notes	<p>Additional Information: Antibody purity is &gt; 95% based on SDS-PAGE. Affinity purified using solid phase Rabbit IgG. Antibody is supplied in 30 mM Triethanolamine, pH 7.2, 5 mM Magnesium Chloride, 0.1 mM Zinc Chloride, 1 % (w/v) BSA, Protease/IgG free. 0.05% (w/v) Sodium Azide is added as a preservative. Based on IEP, this antibody reacts with:</p> <ul style="list-style-type: none"><li>• heavy (γ) chains on rabbit IgG</li><li>• Based on IEP, no reactivity is observed to:</li><li>• non-immunoglobulin rabbit serum proteins</li><li>• light chains on all rabbit immunoglobulins</li><li>• Rabbit IgG F(ab)'2 fragment</li><li>• serum proteins from bovine, horse or human</li></ul> <p>Background: Goat anti-Rabbit IgG Fc, ALP conjugated, min. cross-reactivity to bovine horse or human serum proteins is a secondary antibody, conjugated to ALP, which binds to rabbit IgG (H&amp;L), Fc in immunological assays.</p>