

## **Product Datasheet**

## CHGA - Chromogranin A GRP13222

Species/Host	Chicken
Reactivity	Human
Predicted Reactivity	Human
Immunogen	purified full length native protein UniProt: P10645
Form/Appearance	Liquid in 0.9 % NaCl, 0.1 % sodium azide
Storage	Store at 4°C; make aliquots to avoid working with a stock. Please, remember to spin tubes briefly prior to opening them to avoid any losses that might occur from liquid material adhering to the cap or sides of the tubes.
Note	For research use only.
Clonality	Polyclonal
Purity	Total IgY
MW	51 kDa
Uniprot ID	P10645
Dilution Range	1 : 1000-1 : 5000 (ELISA), 1 : 1000-1 : 5000 (WB)
Application Notes	Additional Information: antibodies are specific against chromogranin A when tested in immunoblotting against urine from patients with carcinoid tumours. IgY fraction at 12.2 mg/ml. IgY fraction was isolated by a two-step PEG precipitation procedure followed by ammonium sulphate precipitation Background: Chromogranin A (a protein of 457 amino acids which is encoded on chromosome 14) is present in neuroendocrine cells throughout the body, including the neuroendocrine cells of the large and small intestine, adrenal medulla and pancreatic islets. It is an excellent marker for carcinoid tumors, phenochromocytomas, paragangliomas, and other neuroendocrine tumors. Coexpression of chromogranin A and neuron specific enolase (NSE) is common in neuroendocrine neoplasms.