

## Product Datasheet

### p53 antibody GRP30

#### Description

This gene encodes tumor protein p53, which responds to diverse cellular stresses to regulate target genes that induce cell cycle arrest, apoptosis, senescence, DNA repair, or changes in metabolism. p53 protein is expressed at low level in normal cells and at a high level in a variety of transformed cell lines, where it's believed to contribute to transformation and malignancy. p53 is a DNA-binding protein containing transcription activation, DNA-binding, and oligomerization domains. It is postulated to bind to a p53-binding site and activate expression of downstream genes that inhibit growth and/or invasion, and thus function as a tumor suppressor. Mutants of p53 that frequently occur in a number of different human cancers fail to bind the consensus DNA binding site, and hence cause the loss of tumor suppressor activity. Alterations of this gene occur not only as somatic mutations in human malignancies, but also as germline mutations in some cancer-prone families with Li-Fraumeni syndrome. Multiple p53 variants due to alternative promoters and multiple alternative splicing have been found. These variants encode distinct isoforms, which can regulate p53 transcriptional activity. [provided by RefSeq]

#### Species/Host

Rabbit

#### Reactivity

Human, Mouse

#### Conjugation

Unconjugated

#### Tested Applications

ChIP, ICC, IF, IHC-P, IP, WB

#### Immunogen

Recombinant protein encompassing a sequence within the center region of human p53. The exact sequence is proprietary.

#### Form/Appearance

Liquid: 1XPBS, 20% Glycerol (pH7). 0.01% Thimerosal was added as a preservative.

#### Concentration

1 mg/ml

#### Storage

Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.

#### Note

For research use only.

#### Isotype

IgG

#### Clonality


Polyclonal

#### Purity


Purified by antigen-affinity chromatography.

#### Uniprot ID

**P04637**



Immunohistochemical analysis of paraffin-embedded Cal27 xenograft, using p53(GRP482) antibody at 1:500 dilution.



p53 antibody immunoprecipitates p53 protein-DNA complex in ChIP experiments. ChIP Sample: HCT116 whole cell lysate/extract treated with CPT 500nM for 6hr A. 5 µg preimmune rabbit IgG. 5 µg of p53 antibody