In der Stockwiese 26

D-85410 Haag/Amper, Germany

Email: info@grp-ak.de Phone: +49 (0)8167 6335

Product Datasheet

MMP2 antibody GRP34

Description Proteins of the matrix metalloproteinase (MMP) family are

involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. This gene encodes an enzyme which degrades type IV collagen, the major structural component of basement membranes. The enzyme plays a role in endometrial menstrual breakdown, regulation of vascularization and the inflammatory response. Mutations in this gene have been

associated with Winchester syndrome and

Nodulosis-Arthropathy-Osteolysis (NAO) syndrome. Two transcript variants encoding different isoforms have been found for this

gene. [provided by RefSeq]

Species/Host Rabbit

Reactivity Human, Mouse

Conjugation Unconjugated

Tested Applications ICC, IF, IHC-P, IP, WB

Immunogen Recombinant protein encompassing a sequence within the cente

region of human MMP2. The exact sequence is proprietary.

Form/Appearance Liquid: 1XPBS, 1% BSA, 20% Glycerol (pH7). 0.025% ProClin 300

was added as a preservative.

Concentration 0.37 mg/ml

Storage Store as concentrated solution. Centrifuge briefly prior to opening

vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid

multiple freeze-thaw cycles.

Note For research use only.

Isotype IgG

Clonality Polyclonal


Purity Purified by antigen-affinity chromatography.

Uniprot ID P08253


Entrez 4313

Dilution Range WB: 1:500-1:3000,ICC: 1:100-1:1000,IHC-P: 1:100-1:1000,IP:

1:100-1:500



Immunofluorescence analysis of methanol-fixed HeLa, using MMP2(GRP486) antibody at 1:200 dilution.



MMP2 protein at cytosol and nucleus on human breast carcinoma by immunohistochemical analysis.
Sample:
Paraffin-embedded human breast carcinoma.
MMP2 antibody (GRP486) dilution:

1:500.