

Product Datasheet

LOX 1 Polyclonal Antibody GRP451

Description Receptor that mediates the recognition, internalization and degradation of oxidatively modified low density lipoprotein (oxLDL) by vascular endothelial cells. OxLDL is a marker of atherosclerosis that induces vascular endothelial cell activation and dysfunction, resulting in pro-inflammatory responses, pro-oxidative conditions and apoptosis. Its association with oxLDL induces the activation of NF-kappa-B through an increased production of intracellular reactive oxygen and a variety of pro-atherogenic cellular responses including a reduction of nitric oxide (NO) release, monocyte adhesion and apoptosis. In addition to binding oxLDL, it acts as a receptor for the HSP70 protein involved in antigen cross-presentation to naive T-cells in dendritic cells, thereby participating in cell-mediated antigen cross-presentation. Also involved in inflammatory process, by acting as a leukocyte-adhesion molecule at the vascular interface in endotoxin-induced inflammation. Also acts as a receptor for advanced glycation end (AGE) products, activated platelets, monocytes, apoptotic cells and both Gram-negative and Gram-positive bacteria (By similarity).

Species/Host Rabbit
Reactivity Human, Mouse, Rat, Rabbit
Conjugation Unconjugated
Tested Applications IHC-P, WB

Immunogen KLH conjugated synthetic peptide derived from Rabbit LOX-1 (public_immunogen_range: 224-274/274)
Form/Appearance Aqueous buffered solution containing 1% BSA, 50% glycerol and 0.09% sodium azide.

Concentration 1ug/ul
Storage Store at -20°C for 12 months.
Note For research use only.

Isotype IgG
Clonality Polyclonal
Purity Purified by Protein A.

Uniprot ID [Q9XTA8](#)
Entrez [100009322](#)

Dilution Range WB: 1:300-1000, IHC-P: 1:200-400

