German Research Products - GRP GmbH

In der Stockwiese 26

D-85410 Haag/Amper, Germany

Email: info@grp-ak.de Phone: +49 (0)8167 6335

Product Datasheet

TNF alpha antibody GRP45

Description This gene encodes a multifunctional proinflammatory cytokine

that belongs to the tumor necrosis factor (TNF) superfamily. This cytokine is mainly secreted by macrophages. It can bind to, and thus functions through its receptors TNFRSF1A/TNFR1 and TNFRSF1B/TNFBR. This cytokine is involved in the regulation of a wide spectrum of biological processes including cell proliferation, differentiation, apoptosis, lipid metabolism, and coagulation. This cytokine has been implicated in a variety of diseases, including autoimmune diseases, insulin resistance, and cancer. Knockout studies in mice also suggested the neuroprotective function of

this cytokine. [provided by RefSeq]

Species/Host Rabbit

Reactivity Human, Mouse, Rat, Bovine

Conjugation Unconjugated

Tested Applications ICC, IF, IHC-P, WB

Immunogen Full length human TNF alpha Recombinant protein.

Form/Appearance Liquid: 1XPBS, 1% BSA, 20% Glycerol (pH7). 0.025% ProClin 300

was added as a preservative.

Concentration 0.04 mg/ml

Storage Store as concentrated solution. Centrifuge briefly prior to opening

vial. For short-term storage (1-2 weeks), store at 4°C. For

long-term storage, aliquot and store at -20°C or below. Avoid

multiple freeze-thaw cycles.

Note For research use only.

Isotype IgG

Clonality Polyclonal

Purity Purified by antigen-affinity chromatography.

Uniprot ID P01375

Entrez 7124


Dilution Range WB: 1:500-1:3000,ICC: 1:100-1:1000,IHC-P: 1:100-1:1000

Non-transfected (â€") and transfected (+) 293T whole cell extracts (30 ?g) were separated by 12% SDS-PAGE, and the membrane was blotted with TNF alpha antibody (GRP497) diluted at 1:1000. The HRP-conjugated anti-rabbit

IgG

antibody was used to

detect



TNF alpha antibody detects TNF alpha protein by western blot

blot analysis. Whole cell extracts (30 ?g) was separated by

12% SDS-PAGE, and the membrane was blotted with TNF alpha antibody (GRP497) diluted

> at 1:500. The