In der Stockwiese 26

D-85410 Haag/Amper, Germany

Email: info@grp-ak.de Phone: +49 (0)8167 6335

Product Datasheet

ASK1 (Thr845) Polyclonal Antibody GRP502

Description Serine/threonine kinase which acts as an essential component of

the MAP kinase signal transduction pathway. Plays an important role in the cascades of cellular responses evoked by changes in the environment. Mediates signaling for determination of cell fate such as differentiation and survival. Plays a crucial role in the

apoptosis signal transduction pathway through

mitochondria-dependent caspase activation. MAP3K5/ASK1 is required for the innate immune response, which is essential for host defense against a wide range of pathogens. Mediates signal transduction of various stressors like oxidative stress as well as by receptor-mediated inflammatory signals, such as the tumor necrosis factor (TNF) or lipopolysaccharide (LPS). Once activated, acts as an upstream activator of the MKK/JNK signal transduction cascade and the p38 MAPK signal transduction cascade through the phosphorylation and activation of several MAP kinase kinases

like MAP2K4/SEK1, MAP2K3/MKK3, MAP2K6/MKK6 and MAP2K7/MKK7. These MAP2Ks in turn activate p38 MAPKs and c-jun N-terminal kinases (JNKs). Both p38 MAPK and JNKs control

the transcription factors activator protein-1 (AP-1).

Species/Host Rabbit

Reactivity Human, Mouse

Conjugation Unconjugated

Tested Applications IHC-P, WB

Immunogen KLH conjugated synthetic phosphopeptide derived from mouse

ASK1 around the phosphorylation site of Thr845

(public_immunogen_range: 825-875/1380)

Form/Appearance Aqueous buffered solution containing 1% BSA, 50% glycerol and

0.09% sodium azide.

Concentration 1ug/ul

Storage Store at -20°C for 12 months.

Note For research use only.

Isotype IgG


Clonality Polyclonal

Purity Purified by Protein A.

Uniprot ID 035099

Entrez 26408

Dilution Range WB: 1:300-1000, IHC-P: 1:200-400





IHC-P of GRP502