In der Stockwiese 26

D-85410 Haag/Amper, Germany

Email: info@grp-ak.de Phone: +49 (0)8167 6335

Product Datasheet

APG5L Polyclonal Antibody GRP533

Description Involved in autophagic vesicle formation. Conjugation with

ATG12, through a ubiquitin-like conjugating system involving ATG7 as an E1-like activating enzyme and ATG1 as an E2-like conjugating enzyme, is essential for its function. The ATG12-ATG5 conjugate acts as an E3-like enzyme which is required for lipidation of ATG8 family proteins and their association to the vesicle membranes. Involved in mitochondrial quality control after oxidative damage, and in subsequent cellular longevity. The ATG12-ATG5 conjugate also negatively regulates the innate antiviral immune response by blocking the type I IFN production pathway through direct association with RARRES3 and MAVS. Also plays a role in translation or delivery of incoming viral RNA to the translation apparatus. Plays a critical role in multiple aspects of lymphocyte development and is essential for both B and T lymphocyte survival and proliferation. Required for optimal processing and presentation of antigens for MHC II. Involved in the maintenance of axon morphology and membrane structures, as well as in normal adipocyte differentiation. Promotes primary ciliogenesis through removal of OFD1 from centriolar satellites and degradation of IFT2 via the autophagic pathway. May play an important role in the apoptotic process, possibly within the modified cytoskeleton. Its expression is a relatively late event in the apoptotic process, occurring downstream of caspase activity. Plays a crucial role in IFN-gamma-induced autophagic cell death

by interacting with FADD.

Species/Host Rabbit

Reactivity Human, Mouse, Rat

Conjugation Unconjugated

Tested Applications IHC-P, WB

Immunogen KLH conjugated synthetic peptide derived from human APG5L

(public_immunogen_range: 118-158/275)

Form/Appearance Aqueous buffered solution containing 1% BSA, 50% glycerol and

0.09% sodium azide.

Concentration 1ug/ul

Storage Store at -20°C for 12 months.


Note For research use only.


Isotype IgG

Clonality Polyclonal

Purity Purified by Protein A.

Unintot ID 09H1V0





IHC-P of GRP533

German Research Products - GRP GmbH In der Stockwiese 26

D-85410 Haag/Amper, Germany

Email: info@grp-ak.de | Phone: +49 (0)8167 6335