

German Research Products - GRP GmbH

In der Stockwiese 26

D-85410 Haag/Amper, Germany

Email: info@grp-ak.de Phone: +49 (0)8167 6335

Product Datasheet

Histone H2A.XS139ph (phospho Ser139) antibody [GT2311] GRP79

Description Histones are basic nuclear proteins that are responsible for the

nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene encodes a member of the histone H2A family, and generates two transcripts through the use of the conserved stem-loop termination motif, and the polyA addition

motif. [provided by RefSeq]

Species/Host Mouse

Reactivity Human, Mouse, Rat

Conjugation Unconjugated

Tested Applications ICC, IF, IHC-P, IP, WB

Immunogen Carrier-protein conjugated synthetic peptide corresponding to

residues around human Histone H2A.XS139ph (phospho Ser139)

The exact sequence is proprietary.

Form/Appearance Liquid: PBS

Concentration 0.92 mg/ml

Storage Store as concentrated solution. Centrifuge briefly prior to opening

vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid

multiple freeze-thaw cycles.

Note For research use only.

Isotype IgG1

Clonality Monoclonal

Purity Affinity purified by Protein G.


Clone ID GT2311

Uniprot ID P16104

Entrez 3014

Dilution Range WB: 1:500-1:3000,ICC: 1:100-1:1000,IHC-P: 1:100-1:1000,IP:

1:100-1:500



Histone H2A.XS139ph (phospho Ser139) antibody [GT2311] detects Histone

H2A.XS139ph (phospho Ser139) protein at

nucleus on mouse testis by immunohistochemical analysis.
Sample:

Paraffin-embedded mouse testis. Histone H2A.XS139ph (phospho Ser139)

antibody [




German Research Products - GRP GmbH

In der Stockwiese 26

D-85410 Haag/Amper, Germany

Email: info@grp-ak.de Phone: +49 (0)8167 6335



Histone H2A.X antibody detects
H2AFX protein at nuclear by immunofluorescent analysis.
Sample: 10?M Cisplatin treated (right) or untreated (left) HeLa cells were fixed in 4% paraformaldehyde for 15 min.
Green: H2AFX protein stained by Histone H2A.Xantibod