

## **Product Datasheet**

## BRCA1 antibody [17F8] - ChIP grade GRP85

Description	BRCA1-17F8 recognizes full-length BRCA1, a 220kDa nuclear phosphoprotein, and does not recognize the Aexon 11 splice variant. In a high proportion of breast and ovarian cancer cell lines, BRCA1 aberrantly mislocates to the cytoplasm. Its usefulness to monitor functional inactivation of BRCA1 in sporadic breast cancers is under active investigation.	
Species/Host	Mouse	KDa EST HET HAP
Reactivity	Human, Mouse	250 - BRCA1
Conjugation	Unconjugated	130 —
Tested Applications	ChIP, ELISA, ICC, IF, IHC-P, IP, WB	BRCA1 antibody [17F8] - ChIP
Immunogen	Protein fragment expressed in E. coli corresponding to amino acids 762-1315.	grade detects BRCA1 protein by western blot analysis.
Form/Appearance	Liquid: PBS	Various whole cell extracts (30 ?g)
Concentration	1.55 mg/ml	were separated by 5% SDS-PAGE, and blotted with BRCA1
Storage	Store as concentrated solution. Centrifuge briefly prior to openin- vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.	and blotted with BRCAT antibody [17F8] - ChIP grade (GRP537) diluted by 1:500. The HRP-conjugated anti
Note	For research use only.	
Isotype	lgG1	
Clonality	Monoclonal	
Purity	Protein G purified	
Clone ID	17F8	The IHC-P analysis of BRCA1
Uniprot ID	P38398	antibody [17F8] - ChIP grade
Entrez	672	was published by Bernard-Gallon DJ and
Dilution Range	WB: 1:500-1:3000,ICC: 1 ?g/ml	colleagues in the journal Breast Cancer Res in 2001.PMID: 11250747